

## Redox Reaction in the Cyclization-aromatization of (*R*)-(+)-Citronellal with FeCl<sub>3</sub>/Acetic Anhydride

Edy Cahyono<sup>a</sup>, Muchalal<sup>b</sup>, Harno Dwi Pranowo<sup>b</sup>, Triyono<sup>b</sup>

<sup>a</sup> Jurusan Kimia, Universitas Negeri Semarang, Kampus Sekaran Gunungpati, Semarang,  
edkim\_unnes@yahoo.co.id

<sup>b</sup> Jurusan Kimia, Universitas Gadjah Mada, Jl. Sekip Utara, Yogyakarta

### ABSTRACT

The effects of temperature, time of reaction, and the reactants mole ratio on the cyclization-acetylation of (*R*)-(+)-citronellal with FeCl<sub>3</sub>/acetic anhydride were studied. Isopulegyl acetate (IPA) and neoisopulegyl acetate (NIPA) were obtained as the main products at room temperature (mole ratio of citronellal: acetic anhydride: FeCl<sub>3</sub> = 6:12:1). Both products are relatively fixed after the 30th minute. The average concentration of IPA at 30-180 minutes was 44.71%, while the average concentration of NIPA was 28.47%. Increasing temperature (80 °C) and the amount of acetic anhydride (mol ratio 2:6:1) gave *p*-cymene (17.53%) and citronellyl acetate (11.31%). Autoredox reaction on the transformation of citronellal into *p*-cymene and citronellyl acetate was studied with the carbon oxidation number concept.

**Key words:** (*R* )-(+)- citronellal, cyclization-aromatization, FeCl<sub>3</sub>, *p*-cymene, redox